

Nemasys® L

Advanced Biocontrol for Horticulture

For control of the larval stages of black vine weevil (*Otiorhynchus sulcatus*) present in soil or container-grown crops

Contains the beneficial nematode *Steinernema kraussei* (88%) and an inert carrier with water (12%)

Product suitable in organic farming according to RCE n° 834/2007

RECOMMENDED RATES

Application method	Application volume	Dose	Pack size 50 million	Pack size 250 million
Open soil	4 litres/m ²	1,000,000/m ²	50m ²	250m ²
Pots / Containers	4 litres/m ²	500,000/m ²	100m ²	500m ²
Drip irrigation (strawberry growbags)	100 ml/plant	25,000/plant	2,000 plants	10,000 plants

The table below gives volumes of diluted Nemasys® L required for individual pots of different sizes, using the recommended pot rate of 125,000 nematodes / litre of water. Note that the nematode volume per pot (ml/pot) decreases as the pot volume increases. The dose rate is related to the surface area.

Pot volume (L)	Drench (ml) / pot	Nematodes/ pot	Pots treated / 50 million pack	Pots treated / 250 million pack
1	46	5,750	8,708	43,542
2	81	10,125	4,940	24,702
5	166	20,750	2,404	12,020
7.5	226	28,250	1,773	8,863
10	278	34,750	1,437	7,187
12	317	39,625	1,263	6,315
14	353	44,125	1,135	5,673
16	386	48,250	1,036	5,180
18	418	52,250	958	4,788
20	448	56,000	894	4,469
22	476	59,500	841	4,203
25	516	64,500	776	3,879
30	576	72,000	695	3,474
35	629	78,625	636	3,179
40	677	84,625	591	2,954
45	720	90,000	556	2,778

DIRECTIONS FOR USE

- Remove all fine filters (50 mesh or smaller) from the application equipment.
- Partly fill the tank with cool (10-20°C), clean water.
- Empty one tray into a bucket and add a small volume of water to form a thick suspension.
- Stir the contents thoroughly making sure all lumps are fully dispersed before adding more water (normally a minimum of 10 litres).
- Add the nematode suspension to the tank and fill with water up to the required level.
- Start the agitator and keep running until the application is completed. If a propeller is used, ensure it does not run at high revolutions (below 150 rpm).
- Apply dose immediately to the whole crop at the appropriate rate.
- Irrigate immediately after soil application to disperse nematodes into soil.

Knapsack sprayer (20L capacity) application instructions (suitable for 50 million packs only):

Open soil application (50 m2)

- Disperse the entire contents of the pack in a bucket containing 10 litres of water to create a stock solution.
- Stir vigorously, immediately take 1 litre of the stock solution and add to knapsack fitted with a coarse spray nozzle and fill with cool, clean water.
- Apply the suspension to a tenth of the area to be treated. Maintain the nematode suspension agitated as nematodes settle out quickly.
- Repeat steps 2 and 3 until the whole area is treated
- Immediately after application water the treated area thoroughly to ensure nematodes are washed into the soil.

81170860 GB 2103

Dosing unit application:

- Set injection rate on dosing unit (normally 1%).
- Considering the pack size, injection rate, area to treat and recommended rates, calculate number of nematodes and volume of water required for a stock solution.
- Make up the stock solution and mix thoroughly to ensure that the nematodes are well dispersed. NOTE: If the suspension is too viscous add an additional known volume of water and adjust the injection rate accordingly.
- Place inlet tube into the stock solution and start the flow of water, so the dosing unit can start injecting nematodes into the outlet pipe.
- Start application ensuring the correct volume of water is being applied per unit area. Throughout application ensure that the nematode suspension remains agitated.
- Apply evenly to the whole area via lance, boom sprayer, sprinklers or drip irrigation system.
- For container applications, avoid overwatering as nematodes can be lost during pot leaking.

Drip irrigation application:

- Allow enough time for nematodes to reach intended area of crop.
- Inject nematodes in the first half of the irrigation cycle and immediately flush the system with enough water to wash nematodes out of irrigation pipes in the second half.
- If multiple blocks are treated, use short irrigation cycles between blocks to ensure an even application of nematodes. Consider spot treatments of drop off zones at the end of drip lines with a lance or knapsack.
- Position irrigation drippers close to the base of the plant. When more than one dripper is used per pot, these should be distributed evenly around the base of the plant.
- Consult your supplier to determine suitability of your drip irrigation system.

APPLICATION CONDITIONS

For optimum results:

- Treatment should be applied when black vine weevil larvae and pupae are present.
- Soil should be moist and between 5°C and 30°C during application and for at least 2 weeks after.
- Avoid applications in direct sunlight; evening applications are recommended.
- Routinely treat new material coming onto the nursery that may contain black vine weevil larvae.
- If product is applied over foliage, irrigate immediately with water to wash the nematodes into the soil.
- Strawberry application: drench each plant individually, around the base of stems. Avoid run off.

PRECAUTIONS

- Agitate nematodes constantly during application.
- Do not use product if past expiry date or incorrectly stored.
- For information relating to which chemicals are compatible with Nemasys® L ask your distributor.
- Once opened, use each entire tray immediately (do not sub-divide or store opened packs or mixed suspension).
- Nemasys® L persists for approximately four weeks; larvae emerging from eggs laid after this time may not be controlled.
- Do not exceed 300 psi/20 bar/2000 kPa pump pressure.
- Nozzle aperture should be greater than 0.5 mm
- Suitable nozzles should produce a coarse spray at the intended application pressure.

SAFETY PRECAUTION

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.
Read label before use.

STORAGE AND DISPOSAL

Storage: Upon receipt, use immediately or store in a refrigerator at or under 5°C. DO NOT FREEZE. Avoid the constant opening of the fridge to minimize temperature fluctuations and do not overfill to allow the movement of cold air. Use entire contents of pack at one time. Do not split or subdivide individual packs as each contains a measured dose of nematodes. Do not store diluted product. KEEP OUT OF REACH OF CHILDREN. THESE PRODUCTS CONTAIN LIVING ORGANISMS THAT REQUIRE SPECIAL HANDLING AND APPLICATION PROCEDURES. FOLLOW THESE INSTRUCTIONS CAREFULLY TO OBTAIN MAXIMUM ACTIVITY.

Disposal: Product and packaging waste resulting from the use of this product, or unused, outdated product, may be disposed of on site or at any approved waste disposal facility in accordance with local regulations.

WARRANTY DISCLAIMER

BASF plc warrants that this product conforms to the description on the label and is reasonably fit for the purpose stated on the label when used in accordance with the directions. BASF PLC DISCLAIMS ALL OTHER WARRANTIES, EITHER EXPRESS OR IMPLIED INCLUDING THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.

REMEDIES

The remedy for injury or damage for this product is limited to, in BASF plc sole discretion, replacement of the product or refund of the purchase price paid. Under no circumstance shall BASF plc be liable for incidental, consequential or special damages.

LIMITATION OF LIABILITY

No one is authorized to change or supplement, whether verbally or in writing, the statements contained on the label. BASF plc and the seller are not liable for personal injury or property damage or loss, including crop damage or lack of performance, due to circumstances beyond their control. BASF does not accept liability for a product user's failure to comply with the directions for use. Buyer and user assume all risks for these events.

Scan QR code for further information:



www.agricentre.basf.co.uk

BASF plc.,
Agricultural Solutions, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG
Telephone: 0161 475 3000
Emergency Information (24 hour Freephone)
00 49 180 22731 12
Technical Enquiries 0845 602 2553 (office hours)

® = Registered trademark of BASF